

## **GROWER PRACTICES WILL INFLUENCE POSTHARVEST FRUIT QUALITY**

**Mary Lu Arpaia**

*Department of Botany and Plant Sciences, University of California, Riverside, CA 92521-0124*

**Zelda Van Rooyen, J. P. Bower**

*Horticultural Science, School of Agricultural Sciences and Agribusiness, University of Natal, Private Bag X01, Scottsville, 3209, South Africa*

**Peter J. Hofman**

*Maroochy Research Station, Queensland Horticulture Institute, Department of Primary Industries, PO Box 5083, Nambour, 4560, Queensland, Australia*

**Allan B. Woolf**

*HortResearch, Mt. Albert Research Center, Private Bag 92 169, Auckland, New Zealand*

Avocado fruit quality relates to several factors and may mean different things to different steps in the postharvest handling of the fruit. The consumer may perceive quality as it relates to appearance, flavor, and dollar value. The packinghouse handler, wholesale distributor, and retailer, in contrast, may perceive quality as ease of handling, uniformity of packaging, and freedom from physical and physiological defects as well as pathological breakdown. In turn, the grower may perceive quality as the overall freedom from defects, optimum size distribution and overall packout. Little attention, traditionally, has been placed on optimization of fruit quality both at harvest and through handling to the consumer.

This paper will examine the role of the preharvest environment and how it may affect the postharvest life of avocado. Essentially, preharvest factors can influence both the rate of fruit development and maturation. The preharvest environment can also have a physical impact on fruit quality: for example, wind or insect scarring may reduce the packout percentage. Finally and most important, but much more difficult to quantify, are those effects that influence susceptibility to physiological and pathological breakdown in the postharvest environment.

The avocado tree exists in a dynamic environment and therefore can respond to changes to that environment in an interactive fashion. The factors listed in Table 1 are neither all inclusive nor mutually exclusive. Several examples will illustrate the interaction between one or more preharvest factors on postharvest fruit quality.

Table 1. Preharvest factors involved in post-harvest fruit quality.

---

Climate/environment
Rootstock/scion
Planting design
Pruning practices
Pest management
Irrigation
Plant growth regulators
Plant nutrition

---

### ENVIRONMENT

Environmental factors include climate (temperature, wind and rainfall), air quality, and positional effects both within a planting and within the tree. Elements such as wind, heavy precipitation, and frost may result in direct loss of the fruit from the postharvest chain due to fruit scarring; increased incidence of plant pathogens associated with high rainfall, especially during flowering (i.e., anthracnose); and loss of fruit related to freeze damage. Temperature during fruit growth and maturation may also influence fruit quality by either hastening or delaying horticultural maturity.

Avocado fruit shape is influenced by growing environment. Fruit grown in cooler environments tend to be more rounded as compared to fruit grown in warmer conditions which tend to be more elongate. In California we have noted that the 'Harvest' (N4 (-) 5) variety (a recent release from the UC Breeding program) will be almost completely round when grown in the relatively cool coastal environment of southern California. Contrasted with this, is the much more elongated fruit shape one observes when the variety is grown in the hot San Joaquin Valley of California. Fruit shape is more tear-dropped shape in intermediate environments. The effect of climate on fruit shape is also evident when examining fruit shape in relation to bloom/fruit set in a given tree. Fruit resulting from "off-bloom" are typically more round than their counterparts set during the main flowering period.

In South Africa the incidence of grey pulp or mesocarp discoloration following storage changes with time between growing districts and time within a harvesting season (Rowell, 1988; Kruger, F.J. and Kritzing, M., 1999; Dixon *et al.*, 2003).

After mild freezes in California, avocado (*Persea americana* Mill.) fruit, although exhibiting no

external damage, may become more susceptible to decay, weight loss, and chilling injury following storage. Relative susceptibility can be correlated to pedicel appearance following exposure to temperatures below -1°C in the field. This characteristic can be used as a marker of fruit quality at harvest following mild freezes (Arpaia, unpublished data). Conversely, Woolf *et al* (1999) demonstrated that exposed sun fruit which are exposed to higher temperatures have a higher tolerance to postharvest heat (50°C hot water treatments) and cold (0°C storage). Several researchers have illustrated the impact of inoculum load and rainfall on the subsequent severity of postharvest decay (Smilanick and Margosan, 2002; Everett *et al.*, 2003, Pak *et al.*, 2003).

### **ROOTSTOCK/SCION AND POLLINIZER EFFECTS**

Kremer-Köhne and Köhne (1992) illustrated the influence of 'Fuerte' vs. 'Hass' avocado on susceptibility to chilling injury. They observed that 'Hass' had a higher percentage of fruit free from physiological disorders following storage than did 'Fuerte'. The use of clonal or "copy" rootstocks for avocado is a relatively recent development (Brokaw, 1987) and subsequently the influence of rootstock on postharvest quality is poorly understood. In spite of this, Marques (2002) demonstrated that rootstock can also influence the incidence and severity of postharvest disease. He demonstrated that 'Hass' trees grown on Velvick clonal rootstock had a lower incidence of postharvest decay as compared to 'Hass' grown on the Duke 7 rootstock under Australian conditions. The work of Degani *et al* (1990) has shown that pollinizer may also influence both fruit size and seed size in 'Fuerte'. More recently (Arpaia, unpublished data) observed that 'Hass' seed length to width ratio was also influenced by the proximity to different pollinizer varieties.

### **PLANTING DESIGN AND PRUNING**

A critical consideration for tropical and subtropical fruits is also the timing of pruning. If pruning or girdling occurs at a time that promotes vegetative growth at the expense of fruit growth, a Ca imbalance and reduction in overall fruit size may occur. Whiley *et al.* (1992) reported higher fruit Ca levels during the first 8 weeks of 'Hass' avocado fruit growth when the spring vegetative flush was controlled with the growth regulator  $\beta$ -[4-chlorophenyl)methyl]- $\alpha$ -(1,1-dimethyl)-1*H*-1,2,4-triazole-1-ethanol (paclobutrazol). Cutting and Bower (1992) demonstrated, under the subtropical growing conditions of South Africa, that avocado fruit borne on trees where vegetative growth was controlled by pruning had higher Ca, Mg, K, and P levels at harvest. Hoffman (un-

published data, 2002) also found that any pruning treatment that stimulates vegetative growth near or during fruit set/growth will reduce quality, and that growth regulators that reduce growth can reduce defects. In Australia, current recommendations it is now believed that canopy management treatments need to avoid increasing vegetative growth during fruit growth otherwise fruit quality will be compromised.

## **PEST MANAGEMENT**

Pest management practices may result in direct fruit loss from scarring and in indirect losses from changes in fruit composition. Thrips scarring, caused by second instar feeding of the avocado thrips on young, developing fruit, is a major concern for growers and is the number one pest controlled in avocados in California. The occurrence of thrips scarring on the fruit results in direct fruit loss as a result of downgrading at the packinghouse. The occurrence of greenhouse thrips in avocado groves also influence not only fruit grade but may also influence the ripening behavior of fruit by slowing or retarding fruit ripening.

## **IRRIGATION**

Irrigation effects on postharvest fruit quality are difficult to quantify. However, Bower (1988) found that preharvest water stress influenced polyphenol oxidase (PPO) levels in ripe avocado fruit after 30 days in storage at 5.5°C. PPO has been associated with mesocarp discoloration in avocados. He also found that preharvest water stress influences the fruit's ability to withstand low O<sub>2</sub> and high CO<sub>2</sub>. Fruit from stressed trees exhibited more physiological disorders following storage and ripening than fruit from nonstressed trees.

## **PLANT GROWTH REGULATORS**

As noted above, the use of plant growth regulators that reduce vegetative vigor may have influence postharvest fruit quality, presumably by influencing the distribution of plant nutrients within the tree. Bower and Cutting (1988) reviewed the literature surrounding endogenous plant growth regulators and their effects on fruit quality. With the exception of ethylene and ABA (via its role in water stress) there is little current evidence that cultural processes greatly influence plant growth regulators and postharvest fruit quality.

## PLANT NUTRITION

The influence of nutritional practices on avocado fruit quality is confusing but overall, research has indicated that the nutritional status of the fruit can impact the postharvest quality of the fruit. Previous work suggested that avocado fruit quality appears to be primarily affected by calcium (Kremer-Köhne *et al.* (1993) and secondarily by nitrogen (Arpaia *et al.*, 1996), boron (Smith *et al.*, 1997), magnesium, potassium (Koen *et al.*, 1990; Witney *et al.*, 1990), and zinc (Vorster and Bezuidenhout, 1988).

Extensive research on 'Fuerte' avocado has been conducted in South Africa. This research focuses on the relationship of postharvest avocado fruit quality and the role of plant nutrition. Witney *et al.* (1990) demonstrated the impact of fruit Ca levels on fruit ripening duration. They reported a significant interaction between these two variables, with high-Ca fruit taking longer to ripen. They also reported on fruit Ca levels during fruit growth from vigorous vs. nonvigorous trees for both 'Hass' and 'Fuerte'. In both instances, fruit borne on nonvigorous trees had higher Ca levels, especially during the early stages of fruit growth. This period also correlates with the time of maximal vegetative flushing. They suggest that it is the early levels of Ca in the fruit that influence subsequent postharvest fruit quality. The work of Whiley *et al.* (1992) confirmed this observation.

Postharvest problems of avocados in South Africa are a combination of several disorders. Swarts (1984) differentiates between pulp spot and mesocarp discoloration or grey pulp. Pulp spot is a blackening of the pulp region surrounding cut vascular bundles and is localized in nature and more prevalent in early season fruit. Mesocarp discoloration or grey pulp is an overall grey-brown flesh discoloration that normally increases with fruit maturity. Vorster and Bezuidenhout (1988) reported lower Zn and Ca levels in fruit exhibiting pulp spot. Smith and Köhne (1992) surveyed a large population of 'Fuerte' avocado trees in which low-yielding trees had poorer fruit quality following storage and lower Ca and Zn levels and higher B levels in flesh tissue. This picture is complicated by the multi-year study of du Plessis and Koen (1992). They concluded that the incidence of grey pulp (or mesocarp discoloration) is strongly correlated with the subsoil Ca and Mg : K ratio. They also reported a significant reduction in the incidence of pulp spot with high levels of subsoil K, which, however, was found to aggravate grey pulp.

An approach recently taken by Van Rooyen and Bower (2004, unpublished data) is to integrate through stepwise multiple regression the nutritional status of the fruit. Their work on 'Pinkerton' avocado demonstrates that excessive nitrogen plays the most significant role in mesocarp discoloration. They also found that decreasing levels of copper, manganese and boron can also

contribute to mesocarp discoloration. Their findings show that the interactions of various plant nutrients can be more important than the individual status of a specific element. For example the nitrogen:calcium ratio was found to significantly influence the development of postharvest disorders whereas the level of calcium by itself did not.

## **POSTHARVEST FACTORS UNDER THE CONTROL OF THE GROWER**

The grower can also play an important role in the postharvest quality of fruit in the manner in which the fruit is handled following harvest prior to delivery to the packinghouse. Eaks (1978) demonstrated that continual temperatures in excess of 25°C following harvest were detrimental to fruit quality from the perspective of fruit ripening. He found that ripening was inhibited when fruit were continuously held at 40°C and abnormal ripening (hard spots) were observed when fruit were held continuously at 30°C. Subsequently Arpaia (1994) observed that cooling delays of 12 or 24 hours when pulp temperature was maintained at 30°C or 40°C during the delay affected ripe fruit quality following storage at 5°C. Woolf *et al.* (1995) observed an increase in avocado tolerance to low temperature following hot air treatments of even 34°C. The rate of ripening was also slowed (decreased shelf life). Such effects have also been noted for fruit exposed to high temperatures (sun exposure) on the tree (Woolf *et al.*, 1999, 2000). It is therefore critical for the grower to protect fruit following harvest from high temperatures. Fruit in the bin can warm rapidly following harvest (Arpaia, unpublished data) reaching temperatures in excess of 25°C within a matter of an hour or two.

## **CONCLUSIONS**

There is ample evidence in current literature that preharvest factors may influence the postharvest fruit quality of avocado. Preharvest factors are perhaps especially critical to the success of handling avocados from Chile since typically long transit times to market are required. Understanding the effects of the preharvest environment on growth and maturation processes and susceptibility to physiological and pathological disorders will help to explain inconsistencies in postharvest fruit performance. There will also be an indirect benefit of this line of research. Typically, growers do not usually understand the postharvest biology of their particular commodity nor do they particularly care since they perceive that “postharvest” is anything past the farm gate. Efforts toward understanding the role of preharvest factors on postharvest quality brings growers into actively controlling the quality of their product and helps to make them willing par-

ticipants in the quest for optimizing product quality.

## LITERATURE CITED

- Arpaia, M. L. 1994. Studies in the postharvest handling of California avocados. *California Avocado Society 1993 Yearbook*. 77:79 – 88.
- Arpaia, M. L., J. L. Meyer, G. W. Witney, G. S. Bender, D. S. Stottlemeyer, and P. R. Robinson. 199). The Cashin Creek nitrogen fertilizer trial – what did we learn? *California Avocado Society Yearbook*, 80:85-98.
- Bower, J.P. 1988. Pre- and postharvest measures for long-term storage of avocados. *South African Avocado Growers' Assn. Yrbk.* 11:68-72.
- Bower, J. P. and J. G. Cutting. 1988. Avocado fruit development and ripening physiology. In: J. Janick (ed.) *Horticultural Reviews*. Volume 10:229-271. Timber Press, Portland, OR.
- Brokaw, W. H. 1987. Field experiences with clonal rootstocks. *South African Avocado Growers' Association Yearbook 1987*. 10:34-36.
- Cutting, J. G. M. and J. P. Bower. 1992. The effect of vegetative pruning on fruit mineral composition and postharvest quality in 'Hass' avocado, p. 403- 407. In: C.J. Lovatt (ed.). *Proc. World Avocado Congr. II*, 21-26 Apr. 1991, Orange, Calif.
- Degani, C., A. Goldring, I. Adato and R. El-Batsri. 1990. Pollen parent effect on outcrossing rate, yield, and fruit characteristics of 'Fuerte' avocado. *HortScience*. 25(4):471-473.
- Dixon, J., H. A. Pak, D. B. Smith, T. A. Elsmly and J. G. M. Cutting. 2003. New Zealand avocado fruit quality: the impact of storage temperature and maturity. *Proceedings V World Avocado Congress (Actas V Congreso Mundial del Aguacate) 2003*. pp. 647-652.
- du Plessis, S. F. and T. J. Koen. 1992. Relationship between mineral nutrition and postharvest fruit disorders of 'Fuerte' avocados, p. 395-402. In: C. J. Lovatt (ed.). *Proc. World Avocado Congr. II*, 21-26 Apr. 1991, Orange, Calif.
- Eaks, I. L. 1978. Ripening, respiration and ethylene production of 'Hass' avocado fruits at 20° to 40°C. *J. Amer. Soc. Hort. Sci.* 103(5):576-578. 1978
- Everett, K. R., J. Rees-George, S. L. Parkes and P. R. Johnston. 2003. Predicting avocado fruit rots by quantifying inoculum potential in the orchard before harvest. *Proceedings V World Avocado Congress (Actas V Congreso Mundial del Aguacate) 2003*. pp. 601-606.

- Koen, T. J., S. F. Du Plessis and J. H. Terblanche. 1990. Nutritional factors involved in physiological post-harvest fruit disorders of avocados (cv. Fuerte). *Acta Horticulturae*, 275:543-550.
- Kremer-Köhne, S. and J. S. Köhne. 1992. Yield and fruit quality of Fuerte and Hass on clonal rootstocks. *South African Avocado Growers' Assn. Yrbk.* 15:69.
- Kremer-Köhne, S., J. S. Köhne and J. M. Schutte. 1993. Effect of potassium, magnesium and nitrogen soil applications on Fuerte avocado fruit quality. *South African Avocado Growers' Association Yearbook*, 16:33-36.
- Kruger, F.J. and Kritzinger, M. 1999. Pinkerton grey pulp problem addressed. *Neltropika*. 306, 5-6.
- Marques, J.R. (2002). 'Hass' avocado fruit quality: the role of fruit minerals and rootstocks. Unpublished PhD thesis, Massey University, New Zealand.
- Pak, H. A., J. Dixon, D. B. Smith, T. A. Elmsly and J. G. M. Cutting. 2003. Impact of rainfall prior to harvest on ripe fruit quality of 'Hass' avocados in New Zealand. *Proceedings V World Avocado Congress (Actas V Congreso Mundial del Aguacate) 2003*. pp. 629-634.
- Rowell, A. W. G. 1988. Cold storage capacity of avocados from different geographic regions. *South African Avocado Growers' Assn. Yrbk.* 11:41-47.
- Smilanick, J., D. and D. Margosan. 2002. Avocado postharvest disease management. *Calif. Avocado Research Symposium*, Oct. 26, 2002.
- Smith, D. G. and J. S. Köhne. 1992. Production potential of Fuerte on seedling rootstocks. *South African Avocado Grower's Assn. Yrbk.* 15:83-85.
- Smith, T. E., C. J. Asher, R. A. Stephenson and S. E. Hetherinton. 1997. Boron deficiency of avocado. 2. Effects on fruit size and ripening In: *Boron in soils and plants* (Bell, R.W. and Rerkasem, B., Eds). Kluwer Academic Publishers, Dordrecht, The Netherlands, 135-137.
- Swarts, D. H. 1984. Post-harvest problems of avocados—Let's talk the same language. *South Africa Avocado Grower's Assn. Yrbk.* 7:15-19.
- Vorster, L. L. and J. J. Bezuidenhout. 1988. Does zinc play a role in reducing pulp spot? *South African Avocado Growers' Assn. Yrbk.* 11:60.
- Wiley, A. W., J. B. Saranah, and B. N. Wolstenholme. 1992. Effect of paclobutrazol bloom sprays on fruit yield and quality of cv. Hass avocado growing in subtropical climates. p. 227-

232. In: C.J. Lovatt (ed.). Proc. World Avocado Congr. II, 21-26 Apr. 1991, Orange, Calif.
- Witney, G. W., P. J. Hofman, and B. N. Wolstenholme. 1990. Effect of cultivar, tree vigour and fruit position on calcium accumulation in avocado fruits. *Sci. Hort.* 44(3-4):269-278.
- Woolf, A.B.; Bowen, J.H.; Ferguson, I.B.; 1999: Preharvest exposure to the sun influences post-harvest responses of 'Hass' avocado fruit. *Postharvest Biology and Technology.* 15(2): 143-153.
- Woolf, A.B.; Watkins, C.B.; Bowen, J.H.; Lay-Yee, M.; Maindonald, J.H.; Ferguson, I.B. 1995: Reducing External Chilling Injury in Stored 'Hass' Avocados with Dry Heat Treatments. *J. Am. Soc Hort. Sci.* 120(6):1050-156.
- Woolf, A.B.; Wexler, A.; Prusky, D.; Kobilier, E.; Lurie, S. 2000: Direct sunlight influences post-harvest temperature responses and ripening of five avocado cultivars. *J. Am. Soc Hort. Sci.* 125(3): 370-376.