

Evaluation of Field Trees for Resistance to *Phytophthora cinnamomi* by Means of the Detached Root Technique

Marius De V. van der Merwe, Erna M.C. Maas, and Johannes M. Kotze

Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria 0002, Republic of South Africa.

Abstract. Avocado trees selected by SAAGA for exceptional growth under apparent root rot pressure were evaluated for resistance by means of the detached root technique. Mycelium fragments of *Phytophthora cinnamomi* were used as an inoculum source instead of zoospores. As means of comparison root segments from root-stocks such as G755, Duke 7 and Edranol with known responses towards *Phytophthora cinnamomi* were used. Nine of the 34 trees evaluated were as resistant as G 755 and six were significantly more resistant than Duke 7. Most of the trees, (i.e. 25) were significantly more resistant than Edranol.

The search for resistance to *Phytophthora cinnamomi* Rands, the causal organism of avocado root rot, was initiated by Dr. G.A. Zentmyer in 1952 (Zentmyer, 1952). According to him, collections have been made in 18 countries and include 15 species of *Persea* and species of other genera in the Lauraceae. Resistance to *P. cinnamomi* in these collections was tested in a nutrient solution test (Zentmyer and Mircetich, 1965), in pots and beds of *P. cinnamomi* infested soil and ultimately in the field (Zentmyer, 1952).

In South Africa, no indigenous *Persea* species occur and the search for resistance is thus restricted to orchard trees showing exceptional signs of vigor under apparent root rot pressure. These trees have been termed "escape" trees.

Obtaining clonal material from these trees for use in resistance tests is a long and tedious procedure. The aim of this study was therefore to evaluate the detached root technique described by Botha, Wehner and Kotze (1989) as a rapid means to assay field trees for resistance to root rot.

Materials and Methods

Roots were obtained from trees selected by Mr. C. Partridge and Mr. D. Westcott of SAAGA. Locality and tree designation are shown in Table 1.

Root tips excised from the different trees were placed separately in plastic containers, filled with moist, sterile vermiculite to prevent dehydration and contamination. The containers were kept in cool boxes, transported to the laboratory and tested within 24 h.

As a control and means of comparison, root tips from two-year-old *P. americana* cultivar Edranol (susceptible) (Snyman *et al.*, 1984) and vegetatively propagated (Frolich and Platt, 1971) *P. americana* selection Duke 7 (moderately tolerant) (Coffey, 1987) as well as *P. schiedeana* Nees selection G755 (tolerant) (Coffey, 1987) were used.

The detached root technique used to test for resistance in avocado rootstocks as described by Dolan and Coffey (1986) and modified by Botha *et al.* (1989) was used. However, inoculum of *P. cinnamomi* consisted of 10 μ L of mycelium suspension. For the mycelium inoculum, 20 5 mm² potato dextrose agar discs (PDA) previously colonized by *P. cinnamomi* were inoculated into 100 mL pea broth prepared as described by Chen and Zentmyer (1970). After shake-incubation at 25C for four days the fungal growth was homogenized for 30 s with an ultra turrax to produce a mycelial suspension.

The excised root tips (ca. 40 mm in length) from each of the different trees, as well as those from the control trees, were placed perpendicularly onto two parallel glass rods in petri dishes containing 15 mL water agar in each as described by Botha *et al.* (1989). Each root tip was inoculated at the region of elongation with 10 μ L of the mycelium homogenate and incubated in the dark at 25C.

Resistance was determined by aseptically cutting the root tips in 4 mm segments after surface disinfecting for 5 s in 70% ethanol. The root segments were then plated out sequentially on PARPH-medium. After incubation at 25C for three days, the segments from which *P. cinnamomi* developed were counted and multiplied by four to obtain the total length of root colonization.

To evaluate whether time after field removal of roots affected expression of resistance, the potted control trees were initially taken to the field and the roots were excised at the same time as those of the field trees. Half of the roots from the control trees were immediately taken to a nearby laboratory and tested as described above. The other half of the roots was kept in the same manner as the root tips of the field trees, until the tests were performed 24 h after detachment.

For all further tests on the field trees, the root tips of the potted control trees remained at the University of Pretoria and were excised at approximately the same time as the root tips of the field trees. The root tips were then kept in the same manner as the root tips of the field trees until the tests could be carried out simultaneously.

Results and Discussion

Time after root detachment (within a 24 h period) did not significantly affect expression of resistance (Table 2). Linear colonization of the excised roots of field trees from each locality as well as the controls are given in Tables 3-6.

Nine of the 34 trees tested showed a higher degree of resistance than G755, although this difference was not significant. Six and 25 trees were found to be significantly more

resistant than Duke 7 and Edranol, respectively. Three of the 34 trees tested were significantly more susceptible than Edranol.

According to Zentmyer and Mircetich (1965) preliminary tests for resistance of rooted cuttings are conducted in a nutrient solution inoculated with *P. cinnamomi*. However, due to the time required to obtain rooted cuttings, an alternative method for evaluating resistance of field trees was investigated. It was found that the detached root technique described by Botha *et al.* (1989) could readily be used when controls such as G755, Duke 7 and Edranol were included in each evaluation. Thus results of this study showed some field trees to be as resistant as the highly acclaimed G755. No previous reports on the "resistant status" of existing avocado trees in South Africa could be found.

We wish to thank Mr. C. Partridge, Mr. D. Westcott and Mr. N. Claassens of SAAGA, for the demarcating of escape trees and assistance.

Literature Cited

- Botha, T., F.C. Wehner, and J.M. Kotze. 1989. An evaluation of in vitro screening techniques for determining tolerance of avocado rootstocks to *Phytophthora cinnamomi*. S. A. Avocado Growers' Assn. Yrbk. 12:60-63.
- Chen, D.W. and G.A. Zentmyer. 1970. Production of sporangia by *Phytophthora cinnamomi* in axenic culture. Mycologia 62:379-402.
- Coffey, M.C. 1987. A look at current avocado rootstocks. California Grower. 11:15-17.
- Dolan, T.E. and M.D. Coffey. 1986. Laboratory screening technique for assessing resistance of four avocado rootstocks to *Phytophthora cinnamomi*. Plant Disease 70:115-118.
- Frolich, E.F. and R.G. Platt. 1971. Use of the etiolation technique in rooting avocado cuttings. Calif. Avocado Soc. Yrbk. 56:97-109.
- Snyman, A.J., C.P. Snyman, and J.M. Kotzé. 1984. Pathogenicity of avocado root rot fungi to Edranol seedlings and Duke 7 rooted cuttings. S. A. Avocado Growers' Assn. Yrbk. 7:80-81.
- Weste, G. 1974. *Phytophthora cinnamomi* - The cause of severe disease in certain nature communities in Victoria. Austral. Journal of Botany 23:67-76.
- Zentmyer, G.A. 1952. Collecting avocados in Central America for disease resistance tests. Calif. Avocado Soc. Yrbk. 32:107-111.
- Zentmyer, G.A. 1984. Avocado Diseases. Trop. Pest Mgmt. 30:388-400.
- Zentmyer, G.A. and S.M. Mircetich. 1965. Testing for resistance of avocado to *Phytophthora* in nutrient solution. Phytopathology 55:487-489.

Table 1. Locality and designation of field trees evaluated for resistance to *P. cinnamomi* by means of the detached root technique.

| Locality | Tree designation |
|-------------|---|
| Agatha | A2, B1, B3, B4, B12, B13, B14 |
| Burgershall | 1D1, 1D2, 1D3, 1D4, 1D5, 1D6, 1D7, 1D8, 1D9, 1D12, 1D13, 1D14 |
| Levubu | 2B1, 2B2, 2B3 |
| Nelspruit | 1Q1, 1Q2, 1Q6, 1Q7, 1Q8, 1 Q9 |
| Tzaneen | Z4 |
| Venda | 1AV, 2AV |
| White River | C1, C2, 1L1 |

Table 2. Linear colonization of potted avocado roots by *P. cinnamomi* at different time intervals after detachment.

| Rootstock | Root colonization | |
|-----------|---------------------------------|----------------------------------|
| | Evaluation 0 h after detachment | Evaluation 24 h after detachment |
| Edranol | 20.45 a ^z | 18.00 a |
| Duke 7 | 6.00 b | 3.40 b |
| G755 | 1.67b | 2.00 b |

^z Values not followed by the same letter differ significantly according to Duncan's multiple range test (P<0.05).

Table 3. Linear colonization of excised root tips of field trees by *P. cinnamomi* after 48 hours; Agatha and Tzaneen sites.

| Rootstock | Linear colonization (mm) of roots |
|-----------|-----------------------------------|
| Edranol | 18.0a ^z |
| B3 | 12.45 ab |
| Z4 | 12.13ab |
| B13 | 11.00 abc |
| B4 | 9.55 bed |
| A2 | 7.27 bed |
| B1 | 6.42 bed |
| B14 | 4.73 bed |
| B12 | 4.44 bed |
| Duke 7 | 3.40 cd |
| G755 | 2.00 d |

^z Values not followed by the same letter differ significantly according to Duncan's multiple range test (P<0.05).

Table 4. Linear colonization of excised root tips of field trees by *P. cinnamomi*

after 48 hours; Nelspruit and White River sites.

| Rootstock | Linear colonization (mm) of roots |
|-----------|-----------------------------------|
| Edranol | 26.08 a ^z |
| 1Q2 | 16.71 be |
| Duke 7 | 14.27 be |
| 1Q7 | 13.83 be |
| 1Q6 | 11.73 bed |
| 1Q9 | 7.72 bed |
| C2 | 5.85 cde |
| G755 | 3.81 cde |
| C1 | 2.00 de |
| 1L1 | 2.00 de |
| 1Q8 | 0.75 e |
| 1Q1 | 0.50 e |

^z Values not followed by the same letter differ significantly according to Duncan's multiple range test (P<0.05).

Table 5. Linear colonization of excised root tips of field trees by *P. cinnamomi* after 48 hours; Levubu and Venda sites.

| Rootstock | Linear colonization (mm) of roots |
|-----------|-----------------------------------|
| Duke 7 | 15.54a ^z |
| 2B3 | 14.84 a |
| 1AV | 14.00 a |
| 2B2 | 13.46 ab |
| Edranol | 13.33ab |
| 2AV | 12.00ab |
| G755 | 10.00 ab |
| 2B1 | 1.54b |

^z Values not followed by the same letter differ significantly according to Duncan's multiple range test (P<0.05).

Table 6. Linear colonization of excised root tips of field trees by *P. cinnamomi* after 48 hours; Burgershall site.

| Rootstock | Linear colonization (mm) of roots |
|-----------|-----------------------------------|
| Edranol | 27.5 a ^z |
| Duke 7 | 21.0ab |
| 1D8 | 20.8 ab |
| 1D12 | 19.3 ab |
| 1D4 | 18.0 ab |
| 1D6 | 15.2 b |
| 1D9 | 14.7 b |
| 1D13 | 12.5 be |
| 1D1 | 12.4 be |
| 1D3 | 11.7 be |
| G755 | 11.5 be |
| 1D5 | 11.4 be |
| 1D2 | 11.3 be |
| 1D14 | 10.6 be |
| 1D7 | 1.5 c |

^z Values not followed by the same letter differ significantly according to Duncan's multiple range test (P<0.05).