

# Interaction of storage, ethylene and ethylene inhibitors on post-harvest quality of 'Maluma'

AA Ernst

Allesbeste Nursery  
PO Box 91, Tzaneen 0850, South Africa  
Tel: +27 15 307 3076. Fax: +27 15 307 7650  
E-mail: [andre@allesbeste.com](mailto:andre@allesbeste.com) Website: <http://www.allesbeste.com>

## ABSTRACT

'Maluma' is a precocious and productive early bearing 'Hass'-like cultivar, commercially released towards the end of 2007. 'Maluma' currently enjoys full plant protection in South Africa (ZA 20043215) and the USA (US PP21, 099 P3). Plant Breeder's Rights are pending in other avocado producing countries. 'Maluma' is an approved South African export cultivar. The responses of 'Maluma' fruit to different post-harvest (pre-storage) fruit treatments with interactions under storage (6°C) and non-storage (ambient) conditions were examined. The aim of this study was to determine the effect of ethylene on post-harvest fruit decays and physiological disorders. The effective management of the negative effects of ethylene through the use of ethylene inhibitors (2,4-dichlorophenoxy acetic acid (2,4-D) and 1-methylcyclopropene (1-MCP)) was also examined. The fruit were ripened at ambient temperature (20-22°C) while the days to ripen were recorded. Fruit were evaluated for the presence of stem-end rot, body rot (anthracnose), vascular browning, red/pink vascular staining and grey pulp. The impact of storage on the rate of ripening was evident, with the least days to ripen with ethylene treated fruit and the most with the 1-MCP treatment. 1-MCP acts as an effective ethylene inhibitor. Less mature fruit are more prone to fungal diseases. Vascular browning is associated with fruit rots. Stored fruit shows less disease related disorders which relates to the faster ripening of the fruit. The frequency and severity of vascular staining significantly increases amongst fruit treated with ethylene prior to storage. This phenomenon, which is ethylene induced, can be efficiently controlled by a 2,4-D (Deccomone®) stem treatment (dip) during picking, together with a 1-MCP (SmartFresh™) treatment prior to cold storage.

**Key words:** 'Maluma', post-harvest, physiological disorders, ethylene, inhibitor, 2,4-D, 1-MCP, rots, vascular browning, vascular staining, grey pulp

## Interacción del Almacenamiento, Inhibidores de Etileno y Etileno en Calidad Post-cosecha de 'Maluma'

AA Ernst

Vivero Allesbeste  
PO Box 91, Tzaneen 0850, República de Sudáfrica  
Tel: +27 15 307 3076. Fax: +27 15 307 7650  
E-mail: [andre@allesbeste.com](mailto:andre@allesbeste.com) Sitio Web: <http://www.allesbeste.com>

## RESUMEN

'Maluma' es un cultivar precoz y productivo, tipo 'Hass', fructífero, lanzado comercialmente a fines de 2007. 'Maluma' actualmente goza de la completa protección de plantas en Sudáfrica (ZA 20043215) y en EE. UU. (US PP21, 099 P3). Los derechos de productores de plantas están pendientes en otros países productores de palta. 'Maluma' es un cultivar de exportación sudafricano aprobado. Se analizaron las respuestas de la fruta de 'Maluma' a diferentes tratamientos de fruta post-cosecha (pre-almacenamiento) con interacciones bajo condi-



ciones de almacenamiento (6°C) y no almacenamiento (ambiente). El objetivo de este estudio fue determinar el efecto del etileno en los deterioros de la fruta post-cosecha y trastornos fisiológicos. El efectivo manejo de los efectos negativos del etileno a través del uso de inhibidores de etileno (2,4-ácido diclorofenoxiacético (2,4-D) y 1-metilciclopropeno (1-MCP)) también se analizó. La fruta maduró a temperatura ambiente (20-22°C) mientras se registraron los días para la maduración. La fruta se evaluó en presencia de putrefacción de tallos, el cuerpo putrefacciones (antracnosis), dorado vascular, manchas vasculares rojas/ rosadas y pulpa gris. El impacto del almacenamiento en la velocidad de maduración fue evidente, siendo menor la cantidad de días para la maduración con la fruta tratada con etileno y mayor con el tratamiento con 1-MCP. El 1-MCP actúa como un inhibidor efectivo de etileno. La fruta menos madura tiene más tendencia a enfermedades de hongos. El dorado vascular se asocia con la putrefacción de la fruta. La fruta almacenada muestra menos trastornos relacionados con enfermedad que se relacione con la maduración más rápida de la fruta. La frecuencia y gravedad de las manchas vasculares aumenta significativamente en las frutas tratadas con etileno antes del almacenamiento. Este fenómeno, inducido por etileno, puede controlarse en forma eficiente con un tratamiento (inmersión) del tallo en 2,4-D (Deccomone®) durante la recolección, junto con un tratamiento de 1-MCP (SmartFresh™) antes del almacenamiento en frío.

**Palabras clave:** 'Maluma', post-cosecha, trastornos fisiológicos, etileno, inhibidor, 2,4-D, 1-MCP, putrefacción, dorado vascular, manchas vasculares, pulpa gris.

## INTRODUCTION

'Maluma' is a precocious and productive early bearing Hass-like cultivar, commercially released at the VI World Avocado Congress held in Viña del Mar, Chile, during November 2007 (Ernst, 2007). 'Maluma' currently enjoys full plant protection in South Africa (ZA 20043215) and the USA (US PP21, 099 P3). Plant Breeder's Rights are pending in the other avocado producing countries. 'Maluma' is an approved South African export cultivar.

The avocado (*Persea americana* Mill.) is a typical climacteric fruit that exhibits a sharp rise in ethylene production during ripening, which occurs after harvest; mature avocado fruit will not ripen while attached to the tree (Blumenfeld & Gazit, 1974). Rhodes (1981) defines the climacteric as "a period in the ontogeny (morphogenesis) of certain fruits during which a series of biochemical changes is initiated by the autocatalytic production of ethylene, marking the change from growth to senescence and involving an increase in respiration and leading to ripening". Therefore ethylene and its autocatalytic production are of vital importance to normal avocado ripening. Tingwa and Young (1975) postulated that some substance, possibly an anion, acts as a ripening regulator and moves either to or from the fruit pedicel once detached from the tree.

Adato and Gazit (1974) found that the faster avocado fruit loses water after picking, the faster they ripen. The ethylene peak occurs earlier if fruit were allowed to become dehydrated, whereas the reverse is true if harvested fruit were infiltrated with water.

Apart from ethylene, abscisic acid (ABA) appears to be a ripening promoter (Bower, 1985). Lieberman, Baker and Sloger (1977) showed an increase in ethylene and ripening following the application of ABA before the climacteric peak. Bruinsma (1981) considered the likely role of ABA to be the stimulation of ethylene biosynthesis once ripening inhibitors of the tree is no longer available after picking. According to Bower (1985), stress is a major factor affecting ABA levels. Once picked, considerable loss of

water from the avocado fruit by transpiration normally takes place. This could create stress conditions in the fruit, leading to ABA accumulation once the fruit water potential approaches zero turgor. Factors other than water stress can also affect ABA synthesis, the most notable in avocado fruit being temperature.

Hershkovitz, Friedman, Goldschmidt and Pesis (2009) illustrated the inducing effect of ethylene on mesocarp discoloration located at the base of the seed in avocado fruit. 1-Methylcyclopropene (1-MCP), the ethylene inhibitor, effectively prevented mesocarp discoloration. Auxins, cytokinins and gibberellins are also inhibitors of ethylene resulting in fruit ripening (Rhodes, 1981). To increase the days to ripen, with consequential improvement of shelf life, Lemmer *et al.* (2008) recommend a 1-MCP dosage rate of 500 parts per billion (ppb) for open market 'Maluma'.

Abscission is an active developmental process occurring at the abscission zone of the fruit pedicel. It is accepted that the increase in ethylene production in the fruit is followed by increased sensitivity of the target cells in the abscission zone to ethylene, which would lead to abscission. In the event to repress post-harvest decay of citrus, the auxin, 2,4-dichlorophenoxy acetic acid (2,4-D), is used as a post-harvest pack house treatment to retard calyx abscission. Commercially the sodium salt of 2,4-D (Deccomone®) is applied to the fruit in a dip treatment at 500 parts per million (ppm) (Cronjé *et al.*, 2005). Deccomone® is registered for the use on citrus in South Africa at a dosage rate of 2 litres/100 litres water (Vermeulen, Grobler & Van Zyl, 2000).

The changes in colour of the fruit skin, as indicator of ripeness, from green to purple/black, is visually more intense with 'Maluma' compared to 'Hass' (Ernst, 2007). The change in pigment is due to an increase in the anthocyanin concentration in the skin tissue during ripening (3-6 days post-harvest) and is almost entirely due to a single anthocyanin: cyaniding 3-O-glucoside (Cox, McGhie, White & Woolf, 2004).

On examining the interaction of ethylene and temperature on post-harvest quality of 'Hass' avocado,



Arpaia (2006) reported on the incidence of pink vascular staining following ripening at 20°C after pre-storage ethylene treatments, subsequently stored at respectively 4 and 12°C (95% RH) for 0, 4 and 14 days. Pink vascular staining was most prevalent in ethylene treated fruit held at 12°C for 14 days. It has been suggested that the occurrence of this disorder is directly linked to the treatment of ethylene in combination with storage, particularly at intermediate temperatures. In relation to storage duration, fruit which did not receive any ethylene displayed a dramatic reduction in days to ripen. A trend of decreasing decay incidence with storage duration was observed, which is believed to be due to the protracted time of ripening of fruit stored for 0 (zero) days.

The aim of this study is to determine the effect of ethylene on post-harvest fruit decays and physiological disorders and to investigate methods to decrease the incidence thereof.

## MATERIALS AND METHODS

The response of 'Maluma' fruit to different post-harvest (pre-storage) fruit treatments with interactions was examined. The treatments included ethylene and ethylene inhibitors under different storage conditions. Fruit treatment commenced within four hours after picking. Three different experiments were conducted, namely:

### Experiment 1:

The experiment was a 2 x 2 x 2 factorial design, laid out on 30 March 2010 and consisted of the following treatments:

- 2 x storage temperatures: control (0 storage at ambient: 20 to 22°C) and 6°C for 7 days;
- 2 x 1-MCP (SmartFresh™) treatments at 0 and 500 ppb at 7°C for 16 hours;
- 2 x ethylene treatments at 0 and 100 ppm at 20°C for 16 hours prior to storage.

The dry matter content of the fruit was 23% (77% moisture).

### Experiment 2:

The experiment was a 2 x 2 x 2 factorial design, laid out on 2 August 2010 and consisted of the following treatments:

- 2 x storage temperatures: control (0 storage at ambient: 20 to 22°C) and 6°C for 7 days;
- 2 x 2,4-D (Deccomone®) stem dip treatments at 0 and 2 litres/100 litres (2%) water at harvest;
- 2 x ethylene treatments at 0 and 100 ppm at 20°C for 16 hours prior to storage.

The dry matter content of the fruit was 33% (67% moisture).

### Experiment 3:

The experiment was a 2 x 2 x 2 x 2 factorial design, laid out on 9 May 2011 and consisted of the following treatments:

- 2 x storage temperatures: control (0 storage

at ambient: 20 to 22°C) and 6°C for 7 days;

- 2 x 2,4-D (Deccomone®) stem dip treatments at 0 and 2 litres/100 litres (2%) water at harvest;
- 2 x 1-MCP (SmartFresh™) treatments at 0 and 500 ppb at 7°C for 16 hours;
- 2 x ethylene treatments at 0 and 100 ppm at 20°C for 16 hours prior to storage.

The dry matter content of the fruit was 23% (77% moisture).

All fruit were treated with Westfalia Biocote™ (92 ml/1 litre water, stem dip) at harvest and prochloraz (1,8 ml/litre water, fruit dip) during packing. Ripening occurred at ambient temperature (20 to 22°C). Days to ripen for each treatment were recorded. Fruit were evaluated for the presence of stem-end rot, other rot (anthracnose), vascular browning, red/pink vascular staining and grey pulp. Data was statistically analysed with the Chi-square % defective test through the aid of the computer software package Minitab 16.

## RESULTS AND DISCUSSION

The results as illustrated in Figures 1 to 5 are from Experiment 1: 'Maluma' typically displays the characteristics of a climacteric fruit as described by Blumenfeld and Gazit (1974). 'Maluma' exhibits a strong autocatalytic production of ethylene which is clearly illustrated (Figure 1) by the days to ripen of the control compared to stored (6°C) fruit, in the absence of applied ethylene, where a substantial decrease in the average time to ripen was visible. This is supported by the fact that the experiment was conducted with fruit only reaching the onset of maturity at a 23% dry matter level.

The shortest days to ripen with the least difference between the control and stored fruit were evident with the ethylene treated fruit (Figure 3). As suggested by Lemmer *et al.* (2008), 1-MCP (500 ppb) effectively increased the average days to ripen of stored and non-stored fruit (Figure 2) and also acts as an effective ethylene inhibitor, resulting in comparable average days to ripen if compared with non-ethylene treated fruit (Figure 4). The dramatic




Figure 1. The effect of storage (7 days at 6°C) on days to ripen of 'Maluma'.



reduction in days to ripen of stored fruit, which did not receive any ethylene, correlates with the findings of Arpaia (2006).

As far as the different fruit rots (stem-end and body rot) and grey pulp are concerned, no significant statistical difference could be found between the different treatments for a particular variable. However, a clear tendency towards a lower percentage of rot is visible among the stored fruit compared with the control (Figures 5 and 7). This supports the observations of Arpaia (2006) who reported on a trend of decreasing decay incidence with an increase in storage duration. It is explained by the protracted time of ripening of fruit stored for 0 (zero) days. Figures 5 and 7 respectively illustrates a typical decrease in the occurrence of stem-end rot and an increase in body rot (mainly anthracnose) associated with the increase in dry matter content of the fruit from 23 to 33% (moisture decrease from 77 to 67%).

The incidence of grey pulp can be explained due to the fact that the fruit collected for the experiment was from young trees (3,5 years old) with relatively

high nitrogen levels. 1-MCP effectively suppressed grey pulp under storage conditions, even with ethylene pre-storage treated fruit (Figure 5). As illustrated in Figures 7 and 9, grey pulp is of no concern in older orchards (9 and 10 years old) where the nitrogen and calcium levels are within the prescribed norms. Ac-




Figure 3. The effect of ethylene (100 ppm) on days to ripen of stored (7 days at 6°C) and non-stored 'Maluma'.




Figure 2. The effect of 1-MCP (500 ppb; SmartFresh) on days to ripen of stored (7 days at 6°C) and non-stored 'Maluma'.




Figure 4. The effect of 1-MCP (500 ppb) and ethylene (100 ppm) on the days to ripen of stored (7 days at 6°C) and non-stored 'Maluma'.




Figure 5. The effect of 1-MCP (500 ppb) and ethylene (100 ppm), individually and in combination, on the intensity of different disorders after ripening of stored (7 days at 6°C) and non-stored 'Maluma'. All values with different superscript letters are significantly different ( $P \leq 0,05$ ).



According to Figure 5 a statistically significant difference ( $P \leq 0,05$ ) exists between the ethylene treated and non-treated fruit with regards to red/pink vascular staining (Figure 6), with the highest incidence among the pre-stored ethylene (100 ppm) treated fruit stored for 7 days at 6°C and ripened at 20°C. 1-MCP (500 ppb) significantly suppressed vascular staining of ethylene treated and stored 'Maluma' fruit, to a level similar to that of non-ethylene treated fruit. The inhibition of ethylene with 1-MCP confirms the findings of Hershkovitz *et al.* (2009). The inducing effect of ethylene on the occurrence of this phenomenon supports the findings of Arpaia (2006) who suggest-



Figure 6. Incidence of pink/red vascular staining following ripening at 20°C after pre-storage (7 days at 6°C) ethylene (100 ppm) treatment of 'Maluma'.




Figure 7. The effect of 2,4-D (2%) and Ethylene (100 ppm), individually and in combination, on the intensity of different disorders after ripening of stored (7 days at 6°C) and non-stored 'Maluma' fruit. All values with different superscript letters are significantly different ( $P \leq 0,05$ ).




Figure 8. The effect of ethylene and ethylene inhibitors on vascular staining of stored (7 days at 6°C) and subsequently ripened 'Maluma' fruit. All values with different superscript letters are significantly different ( $P \leq 0,05$ ).



ed that this disorder is directly linked to the treatment of ethylene in combination with storage, particularly at intermediate temperatures.

In Figure 7 (Experiment 2), 'Maluma' displays a visible lower intensity of stem-end rot and vascular browning at a dry matter content of 33%. Manageable levels of anthracnose (body rot) are observed. 2,4-D effectively reduces the incidence level of vascular staining to zero with non-ethylene treated fruit. With pre-stored ethylene treated fruit, 2,4-D as a stem dip treatment during harvest successfully inhibits ethylene production, resulting in a significant reduction (55%) of vascular staining. In Figure 8 (Experiment 3) this inhibition was not significant; however a clear tendency of ethylene inhibition was illustrated. From the results it is clear that the suppressive effect of 2,4-D is not sustainable during storage, however, it effectively reduces pre-storage ethylene by inhibiting post-harvest stress related ethylene production. During storage 1-MCP effectively and significantly inhibits the effect of pre-storage exposure to ethylene, while in combination with 2,4-D it significantly and fully prevents the development of red/pink vascular staining.

As far as fungal disorders are concerned, no significant difference existed between the different treatments of a particular disorder except for vascular browning, where the stored fruit which were pre-treated with ethylene and 1-MCP had a significantly higher incidence. In all instances the intensity of the disorder was the lowest with the shortest days to ripen. The combination of 1-MCP and 2,4-D effectively reduced the intensity of the particular fungal disorder (Figure 9).

## CONCLUSIONS

'Maluma' is a typical climacteric fruit displaying a sharp rise in ethylene production during ripening, which occurs directly after harvest. The autocatalytic production of ethylene is evident as a sharp decrease in days to ripen occurs with stored fruit, even in the absence of applied ethylene. The dramatic reduction in days to ripen of stored fruit which did not receive any ethylene correlates with the findings of Arpaia (2006).

A clear tendency towards a lower percentage of rots is visible among stored fruit compared with non-stored fruit. This supports the observations of Arpaia (2006) who reported a trend of decreasing decay incidence with an increase in storage duration. It is explained by the protracted time of ripening of non-stored fruit. A decrease in the occurrence of stem-end rot together with an increase in body rot (mainly anthracnose) is typically associated with an increase in the dry matter content of the fruit.

As suggested by Lemmer *et al.* (2008) 1-MCP (500 ppb) effectively increased the average days to ripen of stored fruit and acts as an effective ethylene inhibitor. 1-MCP effectively suppressed grey pulp. Grey pulp with 'Maluma' is of no post-harvest concern, particularly in older orchards where the nitrogen and calcium levels are within the prescribed norms.

The changes in colour of the fruit skin, as indicator of ripeness, from green to purple/black, is visually more intense with 'Maluma' compared to 'Hass' (Ernst, 2007). The change in pigment is due to an increase in the anthocyanin concentration in the skin tissue during ripening (3-6 days post-harvest) (Cox *et al.*, 2004). The red/pink vascular staining




Figure 9. The effect of ethylene and ethylene inhibitors on fungal disorders of stored (7 days at 6°C) and subsequently ripened 'Maluma' fruit. All values with different superscript letters are significantly different ( $P \leq 0,05$ ).



relates to an excessive anthocyanin production as a result of a rapid increase in respiration due to a stress related over exposure to ethylene. To prevent this disorder, it is essential to manage stress related ethylene production with consequential increase in respiration particularly during picking and packing of the fruit.

From the results it is clear that red/pink vascular staining is induced by post-harvest/pre-storage exposure of 'Maluma' fruit to ethylene. The inducing effect of ethylene on the occurrence of this phenomenon supports the findings of Arpaia (2006) who suggested that this disorder is directly linked to the treatment of ethylene in combination with storage, particularly at intermediate temperatures. As an effective ethylene inhibitor 1-MCP (500 ppb) significantly suppressed vascular staining. The inhibition of ethylene with 1-MCP confirms the findings of Hershkovitz *et al.* (2009) regarding the inhibition of mesocarp decolouration with green fruit.

ABA acts as a stimulant for ethylene biosynthesis once ripening inhibitors of the tree are no longer available after picking (Bruinsma, 1981). According to Bower (1985), stress is a major factor affecting ABA levels. Considerable loss of water from the avocado fruit through transpiration, as well as high temperature exposure after picking, creates stress conditions in the fruit, leading to ABA accumulation (Bower, 1985), which in turn stimulates the autocatalytic production of ethylene. As a stem dip treatment during harvest, 2,4-D successfully inhibits ethylene production resulting in a significant reduction of vascular staining. The suppressive effect of 2,4-D is not sustainable during storage, however it effectively reduced pre-storage ethylene by inhibiting post-harvest stress related autocatalytic ethylene. During storage, 1-MCP effectively and significantly inhibits the effect of pre-storage exposure to ethylene, while in combination with 2,4-D it significantly and fully prevents the development of red/pink vascular staining.

It can therefore be concluded that the impact of storage on the rate of ripening is evident, with the least days to ripen with ethylene treated fruit and the most with the 1-MCP treatment. 1-MCP acts as an effective ethylene inhibitor. Less mature fruit are more prone to fungal diseases. Vascular browning is associated with fruit rot. Stored fruit shows less disease related disorders, which relates to the faster ripening of the fruit. The frequency and severity of vascular staining significantly increases amongst fruit treated with ethylene prior to storage. This phenomenon, which is ethylene induced, can be efficiently controlled by a 2,4-D (Deccomo-

ne®) stem treatment (dip) during picking together with a 1-MCP (SmartFresh™) treatment prior to cold storage.

## REFERENCES

- ADATO, I. & GAZIT, S. 1974. Water-deficit stress, ethylene production, and ripening in avocado fruits. *Pl. Physiol.* 53: 45-46.
- ARPAIA, M.L. 2006. *Avocado Postharvest Quality. Continuing Project: Year 7.* Proc Calif Avocado Res Symp, University of California, Riverside. pp. 143-155.
- BLUMENFELD, A. & GAZIT, S. 1974. Development of seeded and seedless avocado fruits. *J. Am. Soc. Hort. Sci.* 9: 442-448.
- BOWER, J.P. 1978. Some aspects of water relations on avocado *Persea americana* (Mill.) Tree and fruit physiology. Pietermaritzburg: MSc. Agric, Thesis, University of Natal.
- BRUINSMA, J. 1981. Hormonal regulation of senescence, ageing, fading and ripening. *Acta Hort.* (138): 141-163.
- COX, K.A., MCGHIE, T.K., WHITE, A. & WOOLF, A.B. 2004. Skin colour and pigment changes during ripening of 'Hass' avocado fruit. *Posth Biol and Tech.* 31: 287-294.
- CRONJÉ, P.J.R., CROUCH, E.M. & HUYSAMER, M. 2005. Postharvest calyx retention of citrus fruit. *Acta Hort. (ISHS)* 682: 369-376.
- ERNST, A.A. 2007. 'Maluma Hass'®: A new released cultivar in comparison with 'Hass'. Proceedings of the VI World Avocado Congress. Viña Dela Mar, Chile.
- HERSHKOVITZ, V., FRIEDMAN, H., GOLDSCHMIDT, E.E. & PESIS, E. 2009. The role of the embryo and ethylene in avocado fruit mesocarp discoloration. *Journal of Experimental Botany*, 60(3): 791-799.
- LEMMER, D., MALUMANE, R. & KRUGER, F.J. 2008. Establishing the effect of smartfresh on five new 'Hass'-like cultivars. ARC-ITSC, Nelspruit.
- LIEBERMAN, M., BAKER, J.E. & SLOGER, M. 1977. Influence of plant hormones on ethylene production in apple, tomato, and avocado slices during maturation and senescence. *Pl. Physiol.* (60): 214-217.
- RHODES, M.J.C. 1981. The maturation and ripening of fruits. In: Thimann, K.V. (ed.) *Senescence in plants.* CRC Press: Boca Raton.
- TINGWA, P.O. & YOUNG, R.E. 1975. Studies on the inhibition of ripening in attached avocado (*Persea americana* Mill.) *Fruits. J. Am. Soc. Hort. Sci.* 100: 447-449.
- VERMEULEN, J.B., GROBLER, H. & VAN ZYL, K. 2000. A guide to the use of plant growth regulants, defoliant and desiccants. NDA, South Africa.

