

## INJECTION OF ESTABLISHED AVOCADO TREES FOR THE EFFECTIVE CONTROL OF PHYTOPHTHORA ROOT ROT

J M DARVAS, J C TOERIEN AND D L MILNE  
WESTFALIA ESTATE, DUIWELSKLOOF

### OPSOMMING

*'n Bale effektiewe wortelvrot beheer is gekry op volwasse avokadobome met die inspuiting van fosetiel-Al, twee maal per seisoen teen 'n dosis van 0,4 g a.b. per m<sup>2</sup> driplyn area. Die behandeling het die voorkoms van **Phytophthora cinnamomi** statisties verminder en dit blyk om 'n veilige kommersiële metode vir die bekamping van wortelvrot te wees, 'n Geringe mate van fitotoksiteit is waargeneem op sommige van die fosetiel-Al ingespuite bome se blare.*

*Metalaksiel ingespuut teen 0,028 g a.b. en Dowco 444 teen 0,4 g a.b. per m<sup>2</sup> driplyn area van die bome het onvoldoende beheer van wortelvrot gegee. Sommige van die bome wat met Dowco 444 ingespuut is het ernstige fitotoksiese simptome gewys.*

### SUMMARY

*Spectacular root rot control has been achieved on fully grown avocado trees by injecting fosetyl-Al twice per season at 0,4 g a.i. per m<sup>2</sup> canopy area. The treatment resulted in a significant reduction of **Phytophthora cinnamomi** incidence in the feeder roots and it has proved to be a safe, commercially applicable technique to combat root rot. A negligible amount of phytotoxicity was observed on the foliage of some of the fosetyl-Al injected trees.*

*Metalaxyl injected at 0,028 g a.i. and Dowco 444 at 0,4 g a.i. per m<sup>2</sup> canopy area of the trees provided an unsatisfactory degree of control against root rot. Some of the trees injected with Dowco 444 showed serious phytotoxic symptoms.*

### INTRODUCTION

Root rot caused by *Phytophthora cinnamomi* Rands is the most devastating disease of avocados (*Persea americana* Mill.) in most avocado producing countries with the result that chemical control of this disease has received much attention. Zentmyer (1955) developed a laboratory method for the screening of fungicides against *P. cinnamomi* and later reported on the controlling effect of Dexon (Zentmyer, 1973). Good progress has been made in the chemical control of root rot with recently developed fungicides of which metalaxyl, fosetyl-Al, Ethazole and Dowco 444 have given encouraging results (Zentmyer, 1980; Darvas, 1983a; Snyman & Kotzé, 1983). The commercial use of metalaxyl and fosetyl-Al has contributed significantly to the

development of the South Africa avocado industry over the past few years.

However, due to the sharp decline in the efficacy of metalaxyl after two years of continued use (Darvas, 1983a) and the slow initial reaction and high application costs with fosetyl-AI (Darvas, 1983a) it was regarded as necessary to develop more effective, low cost chemical control measures. In preliminary trials, three systemic fungicides were tested in the form of a trunk-paint application (Darvas, 1983b). Similar trunk treatments have been investigated on young as well as established trees by Snyman (1982) and Snyman & Kotzé (1983).

This paper reports on the first results obtained from the development of a new technique of injecting systemic fungicides into the trunks of established avocado trees for the control of *Phytophthora* root rot.

## MATERIALS AND METHODS

Ten year old Fuerte trees on root rot susceptible Guatemalan rootstocks were used for the experiment in block 14 of Westfalia Section of Westfalia Estate. The experiment started in 1980 and there were ten single-tree replications in each treatment. Trees were injected in the main trunk by using a slightly modified form of the technique of Buitendag and Bronkhorst (1980). It consisted of drilling holes five mm in diameter and 40 mm deep into the trunk of the trees and using 60 mm capacity syringes. A volume of 20 ml material per syringe was injected under pressure. The number of syringes needed for a tree was established according to tree size based on canopy area ( $m^2$ ) and the injection holes were evenly spaced around the trunk approximately half a meter above ground level. On the completion of the injection, holes were sealed with tree seal. Drills were sterilized after the treatment of each tree by dipping them into 1% Ca-hypochlorite solution to prevent the possibility of other disease transmission. The treatments used in the experiment were as follows:

1. Fosetyl-AI: injected twice per season, first in August and again in October at a rate of 0,4 g a.i. per  $m^2$  canopy area.
2. Metalaxyl: injected at 0,028 g a.i./ $m^2$  in August and in October
3. Dowco 444: injected at 0,4 g a.i./ $m^2$  in August and in October.
4. Control (not injected).

The assessment of the results was based on rating of the trees according to a disease index of 0 (healthy) 10 (dead) during the winter months each year.

Root samples were collected under four trees in each treatment at monthly intervals in the first year of the experiment. A total of 16 freshly killed feeder roots were removed from each sample and surface sterilized in 0,1%  $HgCl_2$  solution for five seconds and rinsed in sterile water. Five 2 mm long pieces were cut from each feeder root and placed onto Potato Dextrose Agar in Petri dishes. The isolated fungi were identified and the mean percentage occurrence of the following organisms was calculated for each treatment: *Phytophthora cinnamomi* Rands, *Pythium* spp., *Cylindrocladium scoparium* Morgan, *Rhizoctonia solani* Kühn, *Macorhominia*

*phaseolina* (Tassi) Goid., *Fusarium oxysporum* Schlecht. and *Cylindrocarpon destructans* (Zins.) Scholten.

TABLE 1: The control of Phytophthora root rot by injecting fungicides into the trunks of established avocado trees.

Treatments	Mean disease rating (0-10) N = 40	
	1980	1981
Fosetyl-A1 injected at 0,4 g a.i./m <sup>2</sup>	5,3	5,0
Metalaxyl injected at 0,028 g a.i./m <sup>2</sup>	4,1	4,6
Dowco 444 injected at 0,4 g a.i./m <sup>2</sup>	4,1	5,0
Control	5,0	4,8

TABLE 2: The incidence of common root pathogens in dead roots of trees injected with fungicides.

Treatments	Mean frequency in dead feeder roots N = 3840						
	<i>P. cinn.</i>	<i>Pythium spp.</i>	<i>C. scop.</i>	<i>R. sol.</i>	<i>M. phaseol.</i>	<i>F. oxysp.</i>	<i>C. destr.</i>
Fosetyl-A1	2,2 b	0,5 a	0,8 a	2,2 a	5,7 a	21,8 a	25,8 a
Metalaxyl	7,1 ab	0,1 a	0,4 a	1,0 a	6,8 a	12,7 a	30,1 a
Dowco 444	6,3 ab	0,07 a	0,7 a	1,0 a	6,6 a	19,8 a	22,2 a
Control	14,1 a	0,7 a	2,7 a	1,2 a	3,6 a	21,2 a	23,2 a

## RESULTS

The mean disease rating of trees in the various treatments at the commencement of the experiment (1980) and in subsequent years is given in Table 1.

The best control of Phytophthora root rot was obtained by injecting fosetyl-AI into the trunk of the trees. The treatment showed little effect in the first year of its use, but it dramatically improved the condition of the trees in the second year. On the other hand, poor control was given by the injection of metalaxyl and Dowco 444 (Table 1.).

The mean percentage incidence of the seven most common avocado root infecting fungi in dead feeder roots in the first year of the experiment is presented in Table 2.

A statistically lower incidence of *P. cinnamomi* was found in roots of fosetyl-AI injected trees than control trees. Metalaxyl and Dowco 444 also tended to reduce the pathogen in the roots, but this reduction was not statistically significant if compared to the control trees. None of the treatments influenced significantly the occurrence of the other pathogens in this survey (Table 2).

A few of the sparsely foliated trees in advanced stages of root rot showed symptoms of phytotoxicity following injections with fosetyl-AI. Symptoms included the development of chlorotic areas between veins of the young leaves. However, these symptoms disappeared in the second year when a large number of new leaves appeared which presumably diluted the fosetyl-AI reaching the foliage. Some of the Dowco 444 injected trees developed phytotoxic symptoms with the formation of abnormally small, brittle leaves often showing tip burn or burn on the edge of the leaf blades and defoliation in the more advanced stages. Trees with less foliage appeared

to be more severely damaged than trees with good leaf cover.

## **DISCUSSION**

The most outstanding control of Phytophthora root rot ever recorded on established avocado trees was achieved by the injection of fosetyl-AI into the trunk of the trees. Various dose rates have been tested in other experiments (unpublished) in which it was confirmed that the rate of 0,4 g a.i. fosetyl-AI per m<sup>2</sup> canopy area gives satisfactory control. The treatment has been tested on Fuerte, Hass, Edranol and Ryan cultivars with good results. As to the after-effect of fosetyl-AI injection, it was found that when trees were cut back eight months after the last injection typical phytotoxic symptoms developed on the leaves of the new flushes. This indicates that the two fosetyl-AI injections in a season give at least a full year's residual effect. Probably there is some residual overlap with the next year's treatment and it may be possible that after the second or third year when trees have regained full vigour and are healthy, fosetyl-AI injected once a year will be sufficient to maintain their condition.

Investigations of the woody parts of the trunk following fosetyl-AI injections revealed that a certain amount of discolouration occurs above and below the point of injection, but this decreases with time and it causes no serious damage to the transporting elements in the trunk (Engelbrecht, 1981). Holes drilled in the trunk callused rapidly after injection, without secondary infections.

The metalaxyl active ingredient for the injection was extracted from the wettable powder formulation and due to its solubility in water, a dose rate of 0,028 g a.i. per m<sup>2</sup> canopy area was administered into the trees. It is possible that a more effective control would be obtained at higher dose rates.

Dowco 444 experimental material at a rate of 0,4 g a.i. per m<sup>2</sup> canopy area exhibited little controlling effect against the disease.

## **ACKNOWLEDGEMENTS**

The technical assistance of Mr D Heyns and MR M Swannack is gratefully acknowledged.

## **LITERATURE CITED**

- BUITENDAG, CH & BRONKHORST, GJ, 1980. Injection of insecticides into tree trunk a possible new method for the control of citrus pests? *Citrus and Subtropical Fruit Journal* No. 556: 5 - 7
- DARVAS, JM, 1983a. Five years of continued chemical control of Phytophthora root rot of avocados. *SAAGA Yearbook* 1983 (in print).
- DARVAS, JM, 1983b. Systemic fungicides applied as trunk paint against root rot of avocados. *SAAGA Yearbook* 1983 (in print).
- ENGELBRECHT, AHP, 1981. Personal communication. Rand Afrikaans University,

Johannesburg.

- SNYMAN, CP, 1982. Die evaluasie van swammiddels teen avokadowortelvrot. *SAAGA Yearbook 1982*: 85 - 93.
- SNYMAN, CP, & KOTZÉ, JM, 1983. Efficacy of systemic fungicides applied as a trunk paint and a sponge band for the control of root rot on five year old avocado trees in the field. *SAAGA Yearbook 1983* (in print).
- ZENTMYER, GA, 1955. A laboratory method for testing soil fungicides with *Phytophthora cinnamomi* as test organism. *Phytopathology* 45: 398 - 404.
- ZENTMYER, GA, 1973. Control of *Phytophthora* root rot of avocados with p-Dimethylaminobenzenediazo Sodium Sulfonate (Dexon). *Phytopathology* 63: 267 - 272.
- ZENTMYER, GA, 1980. *Phytophthora cinnamomi* and the diseases it causes. Monograph No. 10. *Published by the American Phytopathological Society*. 96pp.